











- OSTOVANEH M R, BLACKFORD A, MORAN R A, et al. A novel approach for selecting combination clinical markers of pathology applied to a large retrospective cohort of surgically resected pancreatic cysts[J]. *J Am Med Inform Assoc*, 2017, 24: 145-152.
- [16] HE J, CAMERON J L, AHUJA N, MAKARY M A, HIROSE K, CHOIT M A, et al. Is it necessary to follow patients after resection of a benign pancreatic intraductal papillary mucinous neoplasm?[J]. *J Am Coll Surg*, 2013, 216: 657-665.
- [17] FONG Z V, FERRONE C R, LILLEMOE K D, FERNÁNDEZ-DEL CASTILLO C. Intraductal papillary mucinous neoplasm of the pancreas: current state of the art and ongoing controversies[J]. *Ann Surg*, 2016, 263: 908-917.
- [18] MANDAI K, UNO K, YASUDA K. Does a family history of pancreatic ductal adenocarcinoma and cyst size influence the follow-up strategy for intraductal papillary mucinous neoplasms of the pancreas?[J]. *Pancreas*, 2014, 43: 917-921.
- [19] KAMATA K, KITANO M, KUDO M, SAKAMOTO H, KADOSAKA K, MIYATA T, et al. Value of EUS in early detection of pancreatic ductal adenocarcinomas in patients with intraductal papillary mucinous neoplasms[J]. *Endoscopy*, 2014, 46: 22-29.
- [20] OHTSUKA T, KONO H, TANABE R, NAGAYOSHI Y, MORI Y, SADAKARI Y, et al. Follow-up study after resection of intraductal papillary mucinous neoplasm of the pancreas; special references to the multifocal lesions and development of ductal carcinoma in the remnant pancreas[J]. *Am J Surg*, 2012, 204: 44-48.
- [21] BARTSCH D K, DIETZEL K, BARGELLO M, MATTHAEI E, KLOEPEL G, ESPOSITO I, et al. Multiple small “imaging” branch-duct type intraductal papillary mucinous neoplasms (IPMNs) in familial pancreatic cancer: indicator for concomitant high grade pancreatic intraepithelial neoplasia?[J]. *Fam Cancer*, 2013, 12: 89-96.
- [22] TANAKA M. Current roles of endoscopy in the management of intraductal papillary mucinous neoplasm of the pancreas[J]. *Dig Endosc*, 2015, 27: 450-457.
- [23] AKITA H, TAKEDA Y, HOSHINO H, WADA H, KOBAYASHI S, MARUBASHI S, et al. Mural nodule in branch duct-type intraductal papillary mucinous neoplasms of the pancreas is a marker of malignant transformation and indication for surgery[J]. *Am J Surg*, 2011, 202: 214-219.
- [24] 盛彧,边云,王莉. 壁结节的影像学表现对鉴别胰腺导管内乳头状黏液瘤良恶性的价值[J]. *第二军医大学学报*, 2017, 38: 609-615.
- SHENG Y, BIAN Y, WANG L. Distinguishing benign from malignant intraductal papillary mucinous neoplasms of the pancreas by imaging features of mural nodules[J]. *Acad J Sec Mil Med Univ*, 2017, 38: 609-615.
- [25] JAVIA S, MUNIGALA S, GUHA S, AGARWAL B. EUS morphology is reliable in selecting patients with mucinous pancreatic cyst(s) most likely to benefit from surgical resection[J/OL]. *Gastroenterol Res Pract*, 2017, 2017: 9863952. doi: 10.1155/2017/9863952.
- [26] SPRINGER S, WANG Y, DAL MOLIN M, MASICA D L, JIAO Y, KINDE I, et al. A combination of molecular markers and clinical features improve the classification of pancreatic cysts[J]. *Gastroenterology*, 2015, 149: 1501-1510.
- [27] CARR R A, YIP-SCHNEIDER M T, SIMPSON R E, DOLEJS S, SCHNEIDER J G, WU H, et al. Pancreatic cyst fluid glucose: rapid, inexpensive, and accurate diagnosis of mucinous pancreatic cysts[J]. *Surgery*, 2018, 163: 600-605.
- [28] ARNER D M, CORNING B E, AHMED A M, HO H C, WEINBAUM B J, SIDDIQUI U, et al. Molecular analysis of pancreatic cyst fluid changes clinical management[J]. *Endosc Ultrasound*, 2018, 7: 29-33.
- [29] AL-HADDAD M, DEWITT J, SHERMAN S, SCHMIDT C M, LEBLANC J K, MCHENRY L, et al. Performance characteristics of molecular (DNA) analysis for the diagnosis of mucinous pancreatic cysts[J]. *Gastrointest Endosc*, 2014, 79: 79-87.
- [30] HATA T, DAL MOLIN M, HONG S M, TAMURA K, SUENAGA M, YU J, et al. Predicting the grade of dysplasia of pancreatic cystic neoplasms using cyst fluid DNA methylation markers[J]. *Clin Cancer Res*, 2017, 23: 3935-3944.
- [31] UTOMO W K, LOOIJENGA L H, BRUNO M J, HANSEN B E, GILLIS A, BIERMANN K, et al. A MicroRNA panel in pancreatic cyst fluid for the risk stratification of pancreatic cysts in a prospective cohort[J/OL]. *Mol Ther Nucleic Acids*, 2016, 5: e350. doi: 10.1038/mtna.2016.61.
- [32] LI F, MALLI A, CRUZ-MONSERRATE Z, CONWELL D L, KRISHNA S G. Confocal endomicroscopy and cyst fluid molecular analysis: comprehensive evaluation of pancreatic cysts[J]. *World J Gastrointest Endosc*, 2018, 10: 1-9.
- [33] ROBLES-MEDRANDA C, VARGAS M, OSPINA J, PUGA-TEJADA M, VALERO M, SORIA M, et al. Clinical impact of confocal laser endomicroscopy in the management of gastrointestinal lesions with an uncertain diagnosis[J]. *World J Gastrointest Endosc*, 2017, 9: 389-395.
- [34] BASAR O. Pancreatic cyst biopsy: improvement in diagnosis with micro forceps biopsy[J]. *Cancer Cytopathol*, 2018, 126: 227-228.